

Utah State University

**DigitalCommons@USU**

---

Undergraduate Honors Capstone Projects

Honors Program

---

5-2014

## Dissociation of the Effects of Serotonin Reuptake Inhibitor Fluoxetine in Prelimbic Cortex on Disruption of Timing and Working Memory for Time by Neutral and Negative Emotional Events

Chance Christensen  
*Utah State University*

Follow this and additional works at: <https://digitalcommons.usu.edu/honors>



Part of the [Health and Physical Education Commons](#)

---

### Recommended Citation

Christensen, Chance, "Dissociation of the Effects of Serotonin Reuptake Inhibitor Fluoxetine in Prelimbic Cortex on Disruption of Timing and Working Memory for Time by Neutral and Negative Emotional Events" (2014). *Undergraduate Honors Capstone Projects*. 505.

<https://digitalcommons.usu.edu/honors/505>

This Thesis is brought to you for free and open access by the Honors Program at DigitalCommons@USU. It has been accepted for inclusion in Undergraduate Honors Capstone Projects by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



DISSOCIATION OF THE EFFECTS OF SEROTONIN REUPTAKE INHIBITOR  
FLUOXETINE IN PRELIMBIC CORTEX ON DISRUPTION OF TIMING AND  
WORKING MEMORY FOR TIME BY NEUTRAL AND NEGATIVE EMOTIONAL  
EVENTS

by

Chance Christensen

Thesis submitted in partial fulfillment  
of the requirements for the degree

of

DEPARTMENTAL HONORS

in

Human Movement Science in the Department of Health, Physical Education & Recreation

Approved:

Thesis/Project Advisor  
Eadric Bressel

Departmental Honors Advisor  
Eadric Bressel

~~Director~~ of Honors Program  
Dr. Nicholas Morrison

UTAH STATE UNIVERSITY  
Logan, UT  
Spring 2014

2    **Dissociation of the Effects of Serotonin Reuptake Inhibitor Fluoxetine in Prelimbic Cortex on**  
3    **Disruption of Timing and Working Memory for Time by Neutral and Negative Emotional Events**  
4    Alexander R. Matthews, Chance D. Christensen, Benjamin Zhuoyu Yang, Mona Buhusi and Catalin V  
5    Buhusi\*  
6    USTAR BioInnovation Center, Dept. Psychology, Utah State University, Logan UT 84322  
7    \*Correspondence: Catalin V. Buhusi, USTAR BioInnovations Center, Department of Psychology, Utah  
8    State University, 2810 Old Main Hill, Logan, UT 84322-2810, USA. e-mail: catalin.buhusi@usu.edu

9  
10   Highest degree obtained: ARM, B.S.; CDC, B.S.; BZY, B.A.; MB, Ph.D., M.D.; CVB, Ph.D.

11  
12   Running Title: Fluoxetine effect on interval timing with distracters

13  
14   **Page Limitations**

15 <b>Abstract</b>	<b>244/250</b>
16 <b>Introduction</b>	<b>621/750</b>
17 <b>Methods</b>	<b>994/1500</b>
18 <b>Total</b>	<b>3791/4000</b>

19  
20   **Citations 49/50**

## Abstract

Learning and memory abilities are altered in disorders of the serotonergic system, in disorders such as such as depression, phobias, and post-traumatic stress disorder (PTSD). Among the processes impaired by emotional distracters, and whose dysregulation is documented in affective disorders, is the ability to time in the seconds-to-minutes range, i.e., interval timing. Presentation of distracters during timing tasks result in delays in responding suggesting a failure to maintain subjective time in working memory, as proposed by the Relative Time-Sharing (RTS) model. We investigated the role of the prelimbic cortex in the detrimental effect of anxiety-inducing distracters on the cognitive ability to interval time, using local infusions of selective serotonin reuptake inhibitor (SSRI) fluoxetine in a modified peak-interval procedure with neutral and anxiety-inducing distracters. Given that fluoxetine has beneficial effects on decreasing emotional responses to negative events, we hypothesized that fluoxetine would improve working memory in trials with distracters. Our results revealed a dissociation of the effects of fluoxetine infusion in the prelimbic cortex on interval timing and resource allocation (working memory for time), when neutral and anxiety-inducing distractions occurred. Fluoxetine was effective only during trials with distracters, but not during trials without distracters. Moreover, fluoxetine reduced the harmful effect of the distracters not only when the distracters were anxiety-inducing, but in fact exacerbated their detrimental effects on working memory when they were neutral. Results are discussed in relation to the brain circuits involved in RTS of resources, and the pharmacological management of affective disorders.

Keywords: Interval timing, serotonin, SSRI, fluoxetine, mPFC

## 42 Introduction

43 Learning, memory and emotional processing are dysregulated in affective disorders such as depression,  
44 phobias, and post-traumatic stress disorder (PTSD) resulting in anxiety and altered perception of events  
45 e.g. (Etkin *et al*, 2011; Gil and Droit-Volet, 2009; Hermann *et al*, 2009; Lui *et al*, 2011; Ritchey *et al*,  
46 2011). For example, perception is hindered in individuals with PTSD (Terr, 1983), due to an attentional  
47 bias towards novel or emotionally charged stimuli (Kimble *et al*, 2000). Effective therapies for these  
48 disorders include traditional psychotherapy (Bleiberg and Markowitz, 2005), putative cognitive  
49 enhancers; (Steckler and Risbrough, 2012), and pharmacotherapy (Fletcher *et al*, 2010), prominently  
50 targeting serotonin. Indeed, among the most commonly prescribed antidepressants, 10 out of the top  
51 12 target serotonin (Cipriani *et al*, 2009). In particular, selective serotonin reuptake inhibitors (SSRI) and  
52 putative cognitive enhancers (Floresco and Jentsch, 2011), such as fluoxetine (Prozac®), have beneficial  
53 reductive effects on affective disorder symptoms (Koen and Stein, 2011).

54 One locus of putative serotonergic action is the frontal cortex, responsible for higher-order executive  
55 tasks and functions (Robbins and Arnsten, 2009). Frontal cortex is strongly innervated by serotonin  
56 neurons (Lambe *et al*, 2011) and its damage has severe effects on personality and decision making (Van  
57 Horn *et al*, 2012). Given the prominent expression of serotonergic receptors in the prefrontal cortex  
58 (PFC), a strong modulatory effect by serotonin was ascribed to this region (Puig and Gullledge, 2011).  
59 Therefore, we sought to assess the effects of PFC fluoxetine infusion on cognitive function in the  
60 presence of neutral or negative emotional distracters.

61 Among the processes impaired by emotional distracters, and whose dysregulation is documented in  
62 affective disorders, is the ability to time in the seconds-to-minutes range, i.e., interval timing.  
63 Timekeeping is indeed affected in depression (Droit-Volet, 2013), phobias (Hermann *et al*, 2009),  
64 schizoaffective disorders (Lee *et al*, 2009) and PTSD (Terr, 1983). Presentation of task-irrelevant

distracters during a timing task results in a considerable delay in responding (Buhusi, 2012; Buhusi and Meck, 2009; Matthews *et al*, 2012), suggesting a failure to maintain subjective time in working memory, possibly due to attentional and working memory resources being diverted away from timing (Buhusi *et al*, 2009). Resource re-allocation is exacerbated by anxiety-inducing task-irrelevant distracters, resulting in impairing effects. Most interestingly for the purposes of this paper, antidepressant medication reduces the effect of emotional distracters on interval timing (Matthews *et al*, 2012).

Previous interval timing tasks have shown that the mPFC of is implicated in the ability of distracters to modulate behaviors, and for pharmaceutical manipulations to mediate the negative aspects of the distracter. Matthews *et al* (2012) showed that anxiety-inducing distracters distractibility can be modulated by antidepressant effects focusing on the blockage of dopamine and norepinephrine reuptake within the mPFC of rats. However, serotonin reuptake has also been implicated in anxiety, and is one of the most common forms of pharmaceutical management of anxiety (Cipriani *et al*, 2009). The serotonin transporter (SERT) has been detected in high density within the cortex of rats (Sur *et al*, 1996), and serotonin's post synaptic terminals are also found in great abundance within the mPFC (Puig *et al*, 2011). Unfortunately, the influence of serotonin upon timing with emotional distracters has not been fully elucidated.

We investigated the role of the prelimbic cortex in the detrimental effect of anxiety-inducing task-irrelevant distracters on the cognitive ability to keep track of time, using local infusions of selective serotonin reuptake inhibitor (SSRI) fluoxetine in a modified peak-interval procedure with neutral and anxiety-inducing distracters. Given that fluoxetine has beneficial effects on attention and working memory, e.g., decreasing emotional response to negative events, we hypothesized that fluoxetine would improve maintenance of information in working memory in trials with distracters, resulting in a decrease of the disruptive effect of emotional events on the timekeeping abilities.

## 88    **Methods**

89    *Subjects.* Thirty-three naive 3-4 months old male Sprague-Dawley rats were used. Experimental  
90    procedures were conducted in accordance with the National Institute of Health's Guide for the Care and  
91    Use of Laboratory Animals (2011).

92    *Apparatus.* We used 12 standard rat operant chambers (MED Associates, St. Albans, VT) housed in  
93    sound attenuating cubicles, of which 4 were used for fear conditioning and the other 8 for interval  
94    timing. An 85-dB white noise produced by a white-noise generator was first used during fear  
95    conditioning, and then later used as an anxiety-inducing distracter during the timing task. The  
96    conditioning chambers and the interval timing chambers were made distinctive by using different box  
97    configurations, and different visual, auditory, and odor cues Matthews et al. The fear conditioning  
98    chambers contained a dipper entry space for a liquid dipper (not used). Additionally, no levers were  
99    inserted in the boxes at any time, and no food was ever given. Pine pellets were placed in the waste pan.  
100    In these boxes the grid floor was connected to shockers and scramblers generating a 1-s 0.85-mA foot  
101    shock. In contrast, the interval timing chambers contained two levers situated on the front wall of the  
102    chamber, of which, only the left lever was used. Forty-five mg precision food pellets were delivered as a  
103    reinforcer. The to-be-timed visual stimulus was a 28-V 100-mA house light. A 66-dB background sound  
104    produced by a ventilation fan was present throughout the timing session.

105    *Interval timing.* After autoshaping, rats received fixed Interval (FI) training sessions, during which the  
106    first lever press after 40s was reinforced by food delivery. Afterwards, rats received FI trials randomly  
107    intermixed with non-reinforced peak interval (PI) probe trials. Matthews et al

108    *Fear conditioning.* Rats were randomly assigned into two groups. FEAR rats received 6 pairings of noise  
109    followed by foot shock. NEUTRAL rats were treated identically, but no shock was presented. Rats

received two 30min fear conditioning sessions, one before and one after recovery from surgery, and one fear conditioning session between each testing session.

*Surgery.* Twenty-six-gauge bilateral cannula guides were implanted in aseptic surgery under isoflurane anesthesia into the prelimbic cortex (AP 2.5mm, ML  $\pm$ 0.6mm, DV -3.5mm). Rats were given at least 6 recovery days before resuming timing and conditioning procedures. Rats received 6 PI re-training sessions before PI+N testing.

*Fear test session.* After surgery recovery, FEAR rats received 2 presentations of the noise in extinction followed by 2 noise-shock pairings, at 4min intervals. NEUTRAL rats received no shocks during the fear test. Session lasted 20 minutes.

*Local infusions.* Cannulae injectors aiming at mPFC were lowered into the cannula guides, extending 1mm below the guides. Rats received 0.5 $\mu$ L infusions of either vehicle (n 5% DMSO in saline) or 6 $\mu$ g/side or 0.6 $\mu$ g/side selective serotonin reuptake inhibitor (SSRI) fluoxetine (fluoxetine hydrochloride, Sigma Aldrich, St. Louis, MO), dissolved in vehicle. Infusion order was counterbalanced. See below.

*Timing test sessions with noise and drug infusion.* Rats received local drug infusions followed by a 3hr session of interval timing testing, during which rats received FI and PI trials randomly intermixed with PI trials with noise (PI+N). PI+N trials were similar to PI trials, except that the 5-s white noise was presented (during to-be-timed stimulus), 5s from the onset of the trial. Infusion sessions were separated by 3 retraining sessions.

*Histology.* Rats were anesthetized with isoflurane overdose and transcardially perfused with formalin. Brain sections placed on slides, and stained with crystal violet for histological verification (Figure1). Seven rats were eliminated due to incorrect cannula placement, lost cannulae, failure to acquire the timing task. (NEUTRAL n=14, FEAR n=13)



132 === Figure 1 about here ===

133  
134 *Data Collection and Analysis.* Lever presses during PI and PI+N trials were recorded through a MED  
135 Associates interface. Analyses were conducted on a 100s interval-of-interest starting at the onset of the  
136 to-be-timed signal for all PI trials, and a 20-120s window for PI+N trials. The average response rate was  
137 analyzed as to evaluate: the accuracy of timing (peak time), precision of timing (width of response  
138 function), and peak rate of response. A delay in peak timing in PI+N trials relative to PI trials was  
139 computed and analyzed statistically (Buhusi and Meck, 2000).

140 We further investigated the effect of the presentation of the noise and the effect of the drug on the  
141 dynamics of timing behavior during individual trials. Briefly, during individual trials, the distribution of  
142 lever presses can be approximated by a low-high-low function. Analysis algorithms described in  
143 (Swearingen and Buhusi, 2010) were used to extract the start and stop times. The start time is the time  
144 at which there is a significant increase in response rate during the trial (from the low-to-high states). The  
145 stop time is the point during the trial at which there is a significant decrease in response rate (from the  
146 high-to-low states). PI Trials without temporal control (about 17% [ARM1] of total trials) were eliminated  
147 from individual-trial analyses and there were no exclusion criteria for PI+N trials.

148 Freezing behavior was recorded and scored (in 2.5 s bins) by two independent observers ( $93.24 \pm 1.24\%$   
149 agreement). The percent freezing behavior after the presentation of the noise in extinction was  
150 analyzed to evaluate the fear expression indexed by the rate of decay of freezing behavior after the  
151 presentation of the noise in extinction. Less fear expression was indexed by more negative decay rates  
152 while more fear expression was indexed by more positive decay rates.

153 Timing accuracy (peak time), timing delay after noise, timing precision (width of function), and the start,  
154 stop, and median times were submitted to mixed ANOVAs of peak time, with between-subject variable

group (FEAR, NEUTRAL) and within-subject variables trial type (PI, PI+N) and drug (VEH, FLX 0.6ug, FLX 6ug) were followed by Fisher LSD post-hoc comparisons. To further evaluate the relationship between timing delay and fear expression, the slope of regression lines were calculated using t-tests. Analyses were conducted in SPSS v21 (IBM Corp, Armonk, NY) and Statistica v8 (StatSoft, Tulsa, OK). Statistical tests were evaluated at a significance level of 0.05.

## Results

**No effect of fluoxetine on timing in the absence of distracters.** Rats in both groups' timing peaked about the 40s criterion in PI trials, suggesting that fluoxetine has no effect on interval timing in the absence of distracters (Figure 2). These results were supported by a repeated measures mixed ANOVA on PI times with between-subjects variable (NEUTRAL, FEAR) and within-subject variable drug condition (VEH, FLX 0.6ug, FLX 6ug), which failed to indicate a reliable effect of the drug condition on the average peak time,  $F(2,50) = 2.29$ . These results suggested that fluoxetine has no effect on working memory for time without distraction.

=== Figure 2 about here ===

**Fluoxetine decreases distractibility of negative emotional events, but increases distractibility by neutral events.** In PI+N trials, fluoxetine infusion decreased the delay relative to PI trials by negative emotional events in FEAR rats, but increased distractibility (large delays) by neutral events in NEUTRAL rats. To evaluate these results we computed a delay in peak time in PI+N trials (with noise) relative to PI trials (without noise), by subtracting the estimated peak time in PI trials from the estimated peak time in PI+N trials (Figure 3). A mixed ANOVA of the time delay with between-subjects variable (NEUTRAL, FEAR) and within-subject variable drug condition (VEH, FLX 0.6ug, FLX 6ug) indicated a main effect of group,  $F(1,25)=4.66$ , as well as a group x condition interaction,  $F(2,50)=9.69$ , however failed to find a reliable main effect of drug condition,  $F(2,50)=0.72$ . These results suggested that fluoxetine appeared to have a

178 differential effect upon the distractibility of the groups during the working memory for time task when  
179 emotional distractions were given.

180 === Figure 3 about here ===

181 **No effect of fluoxetine on timing precision or response rate.** Mixed ANOVAs of timing precision (width  
182 of response function) with between-subjects variable (NEUTRAL, FEAR) and within-subject variable drug  
183 condition (VEH, FLX 0.6ug, FLX 6ug) failed to indicate significant effects of group, drug, or interactions,  
184 all  $F_s < 3.09$ . These results suggested that the response curves in PI+N sessions were simply delayed  
185 from PI sessions, indicative of a delay in responding and not an difference in responding placement.

186 **Individual-trial analyses support fluoxetine effects.** Mixed ANOVAs of the start, stop, and median times  
187 in PI trials with between-subjects variable (NEUTRAL, FEAR) and within-subject variable drug condition  
188 (VEH, FLX 0.6ug, FLX 6ug) failed to indicated any differences during PI trials, all  $F_s < 4.0$ . These results  
189 supported the finding that fluoxetine had no effect in PI trials.

190 === Figure 4 about here ===

191 However, significant differences in the start, median, and stop times in the PI+N trials existed supporting  
192 the previous results that suggested that the effect of fluoxetine was only seen during distracted trials.

193 Mixed ANOVAs of the start, stop, and median times indicated main effects of group in the start, median  
194 and stop times, all  $F_s > 5.61$ , and a group x drug interaction in the start and median times,  $F_s > 3.41$ .

195 These results suggest the larger effect of fluoxetine was in the FEAR rats. Also, similarly to what was  
196 found in the curve fitting analyses, the range of the individual trial analyses was not significantly  
197 different in either PI or PI+N trials, supporting the suggestion that the responding curve was simply  
198 delayed.

**Connections between timing delay by fear expression.** Individual animal data was plotted on a regression line for the mean timing delay and fear expression values. The slope of the regression line was compared to null slope, indicating that there was no relationship. T-tests analyses indicated reliable differences from a null slope for the FEAR group under the vehicle condition,  $t(1,12) = 2.53$ , but were not significant under the FLX 6ug condition,  $t(1,12) = 1.62$ . Additionally, the NUETRAL group showed a statistically significant effect under the FLX 6ug condition,  $t(1,13) = 2.51$ ,  $p < 0.05$ , but were not significant under the vehicle condition,  $t(1,13) = 0.77$ . These results supported a paradoxical relationship between the drug conditions and the linear regression slopes.

=== Figure 5 about here ===

## Discussion

Our results revealed a dissociation of the effects of fluoxetine infusion in prelimbic cortex on timing and working memory for time (resource allocation), and between neutral and anxiety-inducing distraction. Fluoxetine was effective in preventing memory disruption by negative emotional distracters, but in fact increased distractibility by neutral distracters. These results are not due to possible changes of the timing mechanisms, as fluoxetine had no effect on timing in the absence of distracters, and had no effect on the precision of timing or on the response rate. These results were confirmed by curve fitting analyses and by analyses of responses in individual-trials.

Similar to what was found in Heilbronner and Meck (2013), the effect of fluoxetine had no impact on non-interrupted (PI) probe trials. Serotonin is thought to have a modulatory effect on interval timing, working memory and action selection (Ho *et al*, 2002; Rogers, 2011). In the past dopamine has been found to be primarily responsible for general timing and implicated in distractibility during trials that have an interruption (Buhusi, 2003; Buhusi and Meck, 2002), as well as an emotional distraction (Matthews *et al*, 2012). As such it is important to see that fluoxetine has no impact upon timing without

222 distraction. Our results indicated that increased serotonin signaling within the mPFC had no impact on  
223 general timing processes.

224 Previous studies have also shown a role of SSRI drug dose on anxiolytic effects (Handley and McBlane,  
225 1993; Pinheiro *et al*, 2007; Salchner and Singewald, 2002; Silva *et al*, 1999). [AM2]Similarly, we saw a dose  
226 dependent effect in the FEAR group, where the delay was decreased as the fluoxetine dose increased.  
227 This result was seen in the timing functions as well as the individual-trial analyses. However, the  
228 anxiogenic result seen in the timing functions was not able to be teased apart in the individual-trial  
229 analyses. This negative result might be due to a lack of power for the analyses. Previous studies have  
230 shown that other analyses have been sensitive enough to find significance in the PI paradigm, in which  
231 individual-trial analyses found no significance (Buhusi and Matthews, 2013).

232 Additionally, the regression lines of the FEAR and NEU groups reversed under the effects of fluoxetine,  
233 which supports the paradoxical effect of fluoxetine seen in the timing functions. It is well known that  
234 one possible acute side effect of SSRIs is anxiogenesis. There is an existing literature showing paradoxical  
235 effects of SSRIs in treatment in both human and animal research (Bagdy *et al*, 2001; Burghardt *et al*,  
236 2004). In human studies, antidepressant drugs have to be administered chronically before therapeutic  
237 relief is seen. However, we were able to see an anxiogenic effect by infusing fluoxetine into the mPFC  
238 directly, which suggests that the effect of serotonin might not have a role in timing, but in retaining  
239 attentional processes within the prefrontal cortex.

240 Previous research has implicated the prefrontal cortex in decision making (Kim *et al*, 2009), including  
241 distraction from memory tasks (Chao and Knight, 1998; Matthews *et al*, 2012). Indeed the prefrontal  
242 cortex of rats has been shown to be analogous to the frontal cortex of nonhuman primates and humans  
243 (Uylings *et al*, 2003; Vertes, 2004). The prefrontal cortex has also been implicated in working memory  
244 for time paradigms, such as the interval timing procedure (Buhusi and Meck, 2005; Kim *et al*, 2009;

245 Matthews *et al*, 2012; Pessoa, 2008). Previous studies have found a high density of serotonin  
246 transporters (SERT) within the cortical areas (Sur *et al*, 1996). SERT is primarily found on the presynaptic  
247 dendrites and its role is to remove serotonin from the synapse. However with fluoxetine, the SERT is  
248 blocked and serotonin is abnormally available within the synapse.

249 Animal[AM3] models of PTSD and anxiety disorders have been developed where an animal is given  
250 classical conditioning where a stimulus is followed by an inescapable footshock (Brown *et al*, 2007;  
251 Matthews *et al*, 2012). Additionally, the time keeping abilities of animals allow scientists to probe  
252 working memory. Over time, animals develop an accurate representation of a time criterion which can  
253 then be used as a probe to identify brain components in working memory tasks when anxiety provoking  
254 situations occur. (Buhusi *et al*, 2005; Matthews *et al*, 2012). During testing sessions, the stimulus was  
255 presented as a task-irrelevant distracter during the timing task resulting in a delay in responding  
256 suggesting a failure to maintain subjective time in working memory, possibly due to attentional and  
257 working memory resources being diverted away from timing, as proposed by the Relative Time-Sharing  
258 (RTS) model (Buhusi, 2003; Buhusi *et al*, 2009).

259 According to the RTS model, both novel and emotional distracters result in attentional processes being  
260 reallocated to the processing of an unexpected stimulus. Previously the attentional processes were  
261 placed onto the primary timing task, however because attentional processes are limited, attention  
262 would be spent on the timing task and upon distracter (Buhusi, 2003; Buhusi *et al*, 2009). However in  
263 disorders such as PTSD, greater attentional emphasis is placed onto stressful distracters, taking more of  
264 the attentional resources available. The RTS model allows us to hypothesize that the distracter should  
265 cause the FEAR group to be shifted in a greater magnitude as compared to the NEU group without any  
266 effect of the drug (VEH condition). However, previous studies have shown that the distractibility of a  
267 stressful stimulus can be partially eliminated using antidepressants within the mPFC of rats (Matthews  
268 *et al*, 2012). The RTS model can explain the paradoxical results. While the distracter is not effective in

269 delaying the timing functions of the NEU group, the NEU group delayed under the effect of FLX,  
270 indicative that increased serotonin redistributed the attentional resources. Conversely, when the  
271 distracter was paired (FEAR) and embedded within the timing task the attentional resources were  
272 redirected upon the noise processing. However, the addition of serotonin might have caused more  
273 resources to be redistributed to processing the timing task. Therefore, we must conclude the serotonin  
274 while having no impact on basal timing, is implicated in the reallocating of attentional processes.

275 In the United States alone, 11% of people over the age of 12 are taking some form of antidepressant  
276 medication (Pratt *et al*, 2011), for depression and anxiety disorders. Proper diagnosis of anxiety  
277 disorders are critical to produce optimal patient outcomes and misdiagnosis and treatment can  
278 exacerbate symptoms. For example, fluoxetine reduces anxiety related symptoms in subjects that  
279 demonstrate anxious behavior (Floresco *et al*, 2011). However, in a population that has no acute  
280 anxiety, fluoxetine treatment has a paradoxical anxiogenic effect (Gorman *et al*, 1987; Pae *et al*, 2004).  
281 Similarly, in our experiment, NEUTRAL rats, which received no noise-shock pairings demonstrated  
282 considerable timing delay in PI+N trials under the high dose of FLX, whereas FEAR rats, which received  
283 noise-shock pairings, were less delayed by the noise. These results suggest that FLX had a differential  
284 effect on the two groups: FLX had an anxiolytic effect when the noise was emotionally charged (FEAR  
285 group), but had an anxiogenic effect when the noise was neutral (NEUTRAL group). Moreover, this may  
286 also explain the individual differences in the effect of FLX on time delay in our experiment (see Figure 5),  
287 as well as individual differences in responsiveness to FLX medication (Oh, 2009) [AM4](Calil, 2001).

288 Fluoxetine is indeed used as an anxiogenic inducer in non-anxious rodents (Bagdy *et al*, 2001), in the  
289 same way that *some* patients experience an exacerbation of anxiety under FLX treatment. Together with  
290 previous studies, our results support a differential effect of SSRI fluoxetine on working memory for  
291 duration related to the context (neutral / fear) in which subjects (patients) are given the treatment.  
292 Further studies are required to elucidate the mechanisms of this paradoxical result.

293     **Funding and Disclosure**

294

295     **Acknowledgments**

296     This research was supported by the National Institutes of Health through grants MH65561 and  
297     MH73057 to CVB. We would like to thank Olivia Hao He and Bogdan Bordeianu for preliminary analyses  
298     and J. Daniel O Bray and Dan Morrison for excellent assistance with animal training. Author contribution:  
299     conceived and designed the experiments: ARM, MB, CVB; performed the experiments: ARM and CDC;  
300     analyzed the data: ARM, CDC, BZY, CVB; wrote the paper: ARM, CDC, MB, CVB.



## 301   References

- 302   Bagdy G, Graf M, Anheuer ZE, Modos EA, Kantor S (2001). Anxiety-like effects induced by acute  
303   fluoxetine, sertraline or m-CPP treatment are reversed by pretreatment with the 5-HT<sub>2C</sub> receptor  
304   antagonist SB-242084 but not the 5-HT<sub>1A</sub> receptor antagonist WAY-100635. *Int J*  
305   *Neuropsychopharmacol* **4**(4): 399-408.
- 306   Bleiberg KL, Markowitz JC (2005). A pilot study of interpersonal psychotherapy for posttraumatic stress  
307   disorder. *Am J Psychiatry* **162**(1): 181-183.
- 309   Brown BL, Richer P, Doyere V (2007). The effect of an intruded event on peak-interval timing in rats:  
310   isolation of a postcue effect. *Behav Processes* **74**(3): 300-310.
- 312   Buhusi CV (ed) (2003). *Dopaminergic mechanisms of interval timing and attention*. CRC Press: Boca  
313   Raton, FL.
- 315   Buhusi CV (2012). Time-sharing in rats: effect of distracter intensity and discriminability. *Journal of*  
316   *Experimental Psychology: Animal Behavioural Processes* **38**(1): 30-39.
- 318   Buhusi CV, Matthews AR (2013). Effect of distracter preexposure on the reset of an internal clock. *Behav*  
319   *Processes*.
- 321   Buhusi CV, Meck WH (2000). Timing for the absence of a stimulus: the gap paradigm reversed. *Journal of*  
322   *Experimental Psychology: Animal Behavioural Processes* **26**(3): 305-322.
- 324   Buhusi CV, Meck WH (2002). Differential effects of methamphetamine and haloperidol on the control of  
325   an internal clock. *Behav Neurosci* **116**(2): 291-297.
- 327   Buhusi CV, Meck WH (2005). What makes us tick? Functional and neural mechanisms of interval timing.  
328   *Nat Rev Neurosci* **6**(10): 755-765.
- 330   Buhusi CV, Meck WH (2009). Relative time sharing: new findings and an extension of the resource  
331   allocation model of temporal processing. *Philos Trans R Soc Lond B Biol Sci* **364**(1525): 1875-1885.
- 333   Burghardt NS, Sullivan GM, McEwen BS, Gorman JM, LeDoux JE (2004). The selective serotonin reuptake  
334   inhibitor citalopram increases fear after acute treatment but reduces fear with chronic treatment: a  
335   comparison with tianeptine. *Biol Psychiatry* **55**(12): 1171-1178.

- 338 Calil HM (2001). Fluoxetine: a suitable long-term treatment. *The Journal of clinical psychiatry* **62 Suppl**  
339 **22**: 24-29.
- 340
- 341 Chao LL, Knight RT (1998). Contribution of human prefrontal cortex to delay performance. *Journal of*  
342 *cognitive neuroscience* **10**(2): 167-177.
- 343
- 344 Cipriani A, Furukawa TA, Salanti G, Geddes JR, Higgins JP, Churchill R, *et al* (2009). Comparative efficacy  
345 and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *Lancet*  
346 **373**(9665): 746-758.
- 347
- 348 Droit-Volet S (2013). Time perception, emotions and mood disorders. *Journal of physiology, Paris* **107**(4):  
349 255-264.
- 350
- 351 Etkin A, Egner T, Kalisch R (2011). Emotional processing in anterior cingulate and medial prefrontal  
352 cortex. *Trends Cogn Sci* **15**(2): 85-93.
- 353
- 354 Fletcher S, Creamer M, Forbes D (2010). Preventing post traumatic stress disorder: are drugs the  
355 answer? *The Australian and New Zealand journal of psychiatry* **44**(12): 1064-1071.
- 356
- 357 Floresco SB, Jentsch JD (2011). Pharmacological enhancement of memory and executive functioning in  
358 laboratory animals. *Neuropsychopharmacology* **36**(1): 227-250.
- 359
- 360 Gil S, Droit-Volet S (2009). Time perception, depression and sadness. *Behav Processes* **80**(2): 169-176.
- 361
- 362 Gorman JM, Liebowitz MR, Fyer AJ, Goetz D, Campeas RB, Fyer MR, *et al* (1987). An open trial of  
363 fluoxetine in the treatment of panic attacks. *Journal of clinical psychopharmacology* **7**(5): 329-332.
- 364
- 365 Handley SL, McBlane JW (1993). Serotonin mechanisms in animal models of anxiety. *Brazilian journal of*  
366 *medical and biological research = Revista brasileira de pesquisas medicas e biologicas / Sociedade*  
367 *Brasileira de Biofisica [et al]* **26**(1): 1-13.
- 368
- 369 Heilbronner SR, Meck WH (2013). Dissociations between interval timing and intertemporal choice  
370 following administration of fluoxetine, cocaine, or methamphetamine. *Behav Processes*.
- 371
- 372 Hermann A, Schafer A, Walter B, Stark R, Vaitl D, Schienle A (2009). Emotion regulation in spider phobia:  
373 role of the medial prefrontal cortex. *Social cognitive and affective neuroscience* **4**(3): 257-267.
- 374
- 375 Ho MY, Velazquez-Martinez DN, Bradshaw CM, Szabadi E (2002). 5-Hydroxytryptamine and interval  
376 timing behaviour. *Pharmacology, biochemistry, and behavior* **71**(4): 773-785.

377  
378 Kim J, Jung AH, Byun J, Jo S, Jung MW (2009). Inactivation of medial prefrontal cortex impairs time  
379 interval discrimination in rats. *Front Behav Neurosci* **3**: 38.

380  
381 Kimble M, Kaloupek D, Kaufman M, Deldin P (2000). Stimulus novelty differentially affects attentional  
382 allocation in PTSD. *Biol Psychiatry* **47**(10): 880-890.

383  
384 Koen N, Stein DJ (2011). Pharmacotherapy of anxiety disorders: a critical review. *Dialogues in clinical*  
385 *neuroscience* **13**(4): 423-437.

386  
387 Lambe EK, Fillman SG, Webster MJ, Shannon Weickert C (2011). Serotonin Receptor Expression in  
388 Human Prefrontal Cortex: Balancing Excitation and Inhibition across Postnatal Development. *PLoS ONE*  
389 **6**(7): e22799.

390  
391 Lee KH, Bhaker RS, Mysore A, Parks RW, Birkett PB, Woodruff PW (2009). Time perception and its  
392 neuropsychological correlates in patients with schizophrenia and in healthy volunteers. *Psychiatry*  
393 *research* **166**(2-3): 174-183.

394  
395 Lui MA, Penney TB, Schirmer A (2011). Emotion effects on timing: attention versus pacemaker accounts.  
396 *PLoS ONE* **6**(7): e21829.

397  
398 Matthews AR, He OH, Buhusi M, Buhusi CV (2012). Dissociation of the role of the prelimbic cortex in  
399 interval timing and resource allocation: beneficial effect of norepinephrine and dopamine reuptake  
400 inhibitor nomifensine on anxiety-inducing distraction. *Front Integr Neurosci* **6**: 111.

401  
402 Pae CU, Lee CU, Lee SJ, Lee C, Paik IH (2004). Fluoxetine-induced mania in an Asian patient. *Psychiatry*  
403 *and clinical neurosciences* **58**(4): 448-449.

404  
405 Pessoa L (2008). On the relationship between emotion and cognition. *Nat Rev Neurosci* **9**(2): 148-158.

406  
407 Pinheiro SH, Zangrossi H, Jr., Del-Ben CM, Graeff FG (2007). Elevated mazes as animal models of anxiety:  
408 effects of serotonergic agents. *Anais da Academia Brasileira de Ciencias* **79**(1): 71-85.

409  
410 Pratt LA, Brody DJ, Gu Q (2011). Antidepressant use in persons aged 12 and over: United States, 2005-  
411 2008. *NCHS data brief*(76): 1-8.

412  
413 Puig MV, Gullledge AT (2011). Serotonin and prefrontal cortex function: neurons, networks, and circuits.  
414 *Molecular neurobiology* **44**(3): 449-464.

415

416 Ritchey M, Dolcos F, Eddington KM, Strauman TJ, Cabeza R (2011). Neural correlates of emotional  
 417 processing in depression: changes with cognitive behavioral therapy and predictors of treatment  
 418 response. *Journal of psychiatric research* **45**(5): 577-587.

419  
 420 Robbins TW, Arnsten AF (2009). The neuropsychopharmacology of fronto-executive function:  
 421 monoaminergic modulation. *Annual review of neuroscience* **32**: 267-287.

422  
 423 Rogers RD (2011). The roles of dopamine and serotonin in decision making: evidence from  
 424 pharmacological experiments in humans. *Neuropsychopharmacology* **36**(1): 114-132.

425  
 426 Salchner P, Singewald N (2002). Neuroanatomical substrates involved in the anxiogenic-like effect of  
 427 acute fluoxetine treatment. *Neuropharmacology* **43**(8): 1238-1248.

428  
 429 Silva MT, Alves CR, Santarem EM (1999). Anxiogenic-like effect of acute and chronic fluoxetine on rats  
 430 tested on the elevated plus-maze. *Brazilian journal of medical and biological research = Revista*  
 431 *brasileira de pesquisas medicas e biologicas / Sociedade Brasileira de Biofisica [et al]* **32**(3): 333-339.

432  
 433 Steckler T, Risbrough V (2012). Pharmacological treatment of PTSD - established and new approaches.  
 434 *Neuropharmacology* **62**(2): 617-627.

435  
 436 Sur C, Betz H, Schloss P (1996). Immunocytochemical detection of the serotonin transporter in rat brain.  
 437 *Neuroscience* **73**(1): 217-231.

438  
 439 Swearingen JE, Buhusi CV (2010). The pattern of responding in the peak-interval procedure with gaps: an  
 440 individual-trials analysis. *Journal of Experimental Psychology: Animal Behavioural Processes* **36**(4): 443-  
 441 455.

442  
 443 Terr LC (1983). Time Sense Following Psychic Trauma. *American Journal of Orthopsychiatry* **53**(2): 244-  
 444 261.

445  
 446 Uylings HB, Groenewegen HJ, Kolb B (2003). Do rats have a prefrontal cortex? *Behav Brain Res* **146**(1-2):  
 447 3-17.

448  
 449 Van Horn JD, Irimia A, Torgerson CM, Chambers MC, Kikinis R, Toga AW (2012). Mapping connectivity  
 450 damage in the case of Phineas Gage. *PLoS ONE* **7**(5): e37454.

451  
 452 Vertes RP (2004). Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse*  
 453 **51**(1): 32-58.

455

456

457 **Figure Legends**

458 Figure 1 Cannulae tip placements. NEUTRAL (n = 14); FEAR (n = 13).

459 Figure 2 Response functions during PI (solid) and noise interrupted PI+N trials (dashed). Vertical lines  
460 indicate estimated peak time in PI (solid) and PI+N trials (dashed). Distance between vertical lines  
461 represents the delay after noise presentation (black rectangle). Vehicle (VEH, left), fluoxetine 0.6ug (FLX  
462 0.6ug, center), and fluoxetine (FLX 6ug, right). \*\*p < 0.01. NEUTRAL (n = 14); FEAR (n = 13).


463 Figure 3 Mean timing delay ( $\pm$  SEM) in PI+N relative to PI trials. Vehicle (VEH, left), fluoxetine 0.6ug (FLX  
464 0.6ug, center), and fluoxetine (FLX 6ug, right). Post-hoc comparisons: \* p < 0.05, \*\* p < 0.01. NEUTRAL (n  
465 = 14); FEAR (n = 13).

466 Figure 4 Mean ( $\pm$  SEM) start (left), median (middle), and stop times (right) in PI+N trials. Post-hoc  
467 comparisons: \* p < 0.05, \*\* p < 0.01. NEUTRAL (n = 14); FEAR (n = 13).


468 Figure 5 Scatterplots of timing delay by fear expression in NEUTRAL (left, n = 14) and FEAR rats (right, n =  
469 13). Vehicle (VEH, dotted lines) and fluoxetine (FLX 6ug, solid lines). Fear expression was indexed by the  
470 rate of decay of freezing behavior after the presentation of the noise in extinction. Less fear expression  
471 (left side of the x-axis) was indexed by more negative decay rates while more fear expression (right side  
472 of the x-axis) was indexed by more positive decay rates.

473 **Figure 1**

474




475





479 Figure 3




480

481

482

483 Figure 4

484



485  
486

